

学校编码: 10384

分类号_____密级_____

学号: 200434037

UDC _____

厦 门 大 学

硕 士 学 位 论 文

源头流域氮的来源迁移与转化

Nitrogen Sources, Transports and Transformations in
Headwater Catchment

王吉苹

指导教师姓名: 曹文志副教授

专 业 名 称: 环境科学

论文提交日期: 2007 年 5 月

论文答辩时间: 2007 年 6 月

学位授予日期: 2007 年 月

答辩委员会主席: 朱鹤健 教授

评 阅 人: 张玉珍 高级工程师

陈松彪 副教授

2007 年 6 月

厦门大学博硕士论文摘要库

厦门大学学位论文原创性声明

兹呈交的学位论文，是本人在导师指导下独立完成的研究成果。
本人在论文写作中参考的其他个人或集体的研究成果，均在文中以明确方式标明。本人依法享有和承担由此论文产生的权利和责任。

声明人（签名）：

年 月 日

厦门大学博硕士论文摘要库

厦门大学学位论文著作权使用声明

本人完全了解厦门大学有关保留、使用学位论文的规定。厦门大学有权保留并向国家主管部门或其指定机构送交论文的纸质版和电子版,有权将学位论文用于非赢利目的的少量复制并允许论文进入学校图书馆被查阅,有权将学位论文的内容编入有关数据库进行检索,有权将学位论文的标题和摘要汇编出版。保密的学位论文在解密后适用本规定。

本学位论文属于

- 1、保密 (), 在 年解密后适用本授权书。
- 2、不保密 (☒)

(请在以上相应括号内打“√”)

作者签名:

日期: 年 月 日

导师签名:

日期: 年 月 日

厦门大学博硕士论文摘要库

摘 要

过量施肥、化石燃料的燃烧等人类活动带来的氮负荷问题对陆地和水生生态系统的影响迅速扩大，引起了全世界的关注。当流域陆地生态体系的氮素饱和后，过量的氮素会通过地表径流和地下潜流排入溪流、湖泊、河流和海洋中。因为溪流直接迁移了流域体系过多的氮素，研究氮在溪流中的循环过程之前了解流域氮素流失到溪流的过程机制至关重要。因此，对五川流域氮素流失的过程机理进行研究以及探讨氮在溪流中的迁移转化行为，有利于阐明流域体系的氮在溪流中的循环过程。

本研究融合多学科知识，在 GIS 技术支持下，综合运用现场定位试验、模型模拟、同位素示踪等多种研究手段与方法，揭示了流域氮的地表流失、地下渗漏淋失等的迁移转化特征并量化，阐明了可溶性无机氮在溪流中的迁移转化过程。主要研究结论如下：

第一，氮的地表径流输出过程及机制。流域氮的地表径流输出主要受降雨、径流、土地利用和施肥状况的多种因素影响，表现出明显的季节性特征。污染物输出主要集中在 4-9 月。2004-2006 年地表径流各形态氮的总输出负荷逐年增加，总氮年输出负荷最小值是 2004 年的 $37.11 \text{ kg N hm}^{-2}$ ，最大值是 2006 年的 108 kg N hm^{-2} 。与五川流域年平均总氮输入 $421.4 \text{ kg N hm}^{-2}$ 相比，地表径流总氮输出平均为 $68.35 \text{ kg N hm}^{-2}$ ，占总输入的 16.2%，其中降雨径流平均输出总氮 $45.51 \text{ kg N hm}^{-2}$ ，占流域总氮输入的 10.8%，基流平均输出总氮 $22.84 \text{ kg N hm}^{-2}$ ，占流域总氮输入的 5.4%。地表径流氮素流失量逐年递增，流失的氮排放到溪流河道中，使溪流氮负荷增加，水质恶化。形态组成上，溶解态氮占 TN 主要比例（DTN/TN=0.84）；在 DTN 中，DIN 占 89.5%，DON 仅占 10.5%；在 DIN 中， $\text{NO}_3\text{-N}$ 占 65.5%， $\text{NH}_4\text{-N}$ 占 34.5%。

第二，氮的淋失过程及机制。五川流域氮的渗漏淋失主要与施肥量、土壤类型、降雨等因素有关。氮淋失在降雨密集的 4-9 月较为严重。 $\text{pH}<5$ 的酸性土壤带正电荷致使铵氮更易向下淋洗造成“富集”，氮淋失以铵氮为主（约 40%）。利用校验后的 GLEAMS 模型，模拟得到 2004-2006 不同土地利用地块硝氮淋失量从 5.00 到 $50.79 \text{ kg N hm}^{-2} \text{ a}^{-1}$ 不等。3 年中，流域加权土地面积总氮年渗漏淋

失负荷范围为 27.47-99.97 kg N hm⁻²，平均渗漏淋失负荷为 69.13 kg N hm⁻²，地下渗漏淋失总氮占流域总氮输入（421.4 kg N hm⁻²）的 16.4%；而 NO₃-N 年渗漏淋失负荷范围为 7.89-28.31 kg N hm⁻²，平均 19.68 kg N hm⁻²，占流域 TN 输入的 4.67%。

第三，五川溪流氮的迁移和转化过程。硝氮与 $\delta^{15}\text{N}$ 、亚硝氮、溶解氧、溶解有机碳的变化关系表明溪流沿程硝化作用是氮在溪流河道的主要生物地球化学过程，也是溪流硝氮浓度沿程升高的主要原因。流域沿程耕地面积比例的增加和面源污染物随地表径流的排入也是溪流沿程铵氮和硝氮浓度升高，特别是铵氮浓度升高的主要原因之一。五川源头溪流较浅的河床为氮的迁移和转化提供了相对较活泼的载体。溪流是联接流域陆地和水体生态系统氮素的重要纽带。

关键词：氮； 来源与转化； 源头流域；

ABSTRACT

Human activities such as fertilizer application, fossil fuel combustion have resulted in increased nitrogen(N) loads for terrestrial and aquatic ecosystems causing worldwilde concern. As terrestrial ecosystems become saturated with N, excess N moves with surface runoff and groundwater flow to streams, lakes, rivers, and coastal oceans. Headwater streams convey water and nutrients to larger streams and despite their relatively small dimensions, play a disproportionately large role in nitrogen transformations on the landscape. Quantitative information on N cycling in streams is needed to understand how N loading from watersheds for streams transporting much of this N directly from terrestrial ecosystems.

In this study, Water quality monitoring, field measurement, mechanism model, isotope tracing method and GIS technique were linked to estimate N flux of streamflow discharge and leaching. The work provides a sound understanding of the influence of surface runoff and groundwater flow on stream N, and highlights transformation and biogeochemistry behavior of dissolved nitrogen in stream.

First, annual and seasonal patterns of N loss in streamflow were evaluated based on monitoring data of water quality and flow in 2004-2006. N loss in stormflow was positively related to the ratio of arable land, and varied greatly among the representative subwatersheds, reflecting the differences in precipitation, land cover, and N inputs. The annual maximum total N export was 108 kg N hm^{-2} of 2006, and the minimum was $37.11 \text{ kg N hm}^{-2}$ of 2004. Compared to average annual N inputs, $421.4 \text{ kg N hm}^{-2}$, average annual total N loss of these three years from streamflow was $68.35 \text{ kg N hm}^{-2}$. Storm runoff conveyed total N $45.51 \text{ kg N hm}^{-2}$, which contributed to 10.8% of total N inputs and baseflow conveyed total N $22.84 \text{ kg N hm}^{-2}$, which contributed to 5.4% of total N inputs. The riverine export of dissolved total nitrogen (DTN), formed 84% of the total flux and the dissolved inorganic nitrogen (DIN) formed 89.5% of DTN. $\text{NO}_3\text{-N}$ and $\text{NH}_4\text{-N}$ were main forms of DIN, contributed to 65.5% and 34.5% respectively.

Second, N leaching in Wuchuan catchment was evaluated using shallow water concentration and GLEAMS (*Groundwater Loading Effects of Agricultural Management Systems*) model. The variation of N concentration in shallow aquifers indicated that higher fertilization under land of

vegetable and commercial crops in spring provided the larger N source. Meanwhile, low hydrologic table in winter (dry season) resulted in a concentrating effect of N concentration. The peak N export in leachate occurred between July to September when rainfall intensity was very high. N leaching was dominated by ammonium form (40%) rather than nitrate form (20%), as a result of acid soil being positively charged with electricity when $\text{pH} < 5$. Nitrate export rate differed substantially with landuse and varied from 5.00 to 50.79 kg N hm^{-2} in 2004-2006, with the area-weighted value of 19.70 kg N hm^{-2} accounting for 4.7% of total N inputs in Wuchuan catchment.

Third, This study found that NH_4 and NO_3 concentrations in streams were in dynamic balance, controlled by input, nitrification, biological uptake, sorption, and regeneration. The relationship between $\text{NO}_3\text{-N}$ with δN^{15} , $\text{NO}_2\text{-N}$, dissolved oxygen, dissolved organic carbon indicated that nitrification was the main biogeochemical process of Wuchuan stream, which played a large role in increasing nitrate levels during stream transport over a relatively short distance. With the area of arable land increment, non-point pollution discharging into the stream was also responsible for $\text{NH}_4\text{-N}$ and $\text{NO}_3\text{-N}$ concentration downstream rising, especially for $\text{NO}_3\text{-N}$. In headwater streams, uptake and removal processes occur mainly on sediments and biofilms covering submerged surfaces. Thus, the shallow depths and high surface-to-volume ratios characteristic of the Wuchuan stream supply a relative reactive channel conduit.

Key Words: Headwater stream; Nitrogen; Sources and transformations;

目 录

表索引	IX
图索引	XI
1 绪论	1
2 国内外研究进展	3
2.1 农田氮素渗漏淋失的研究进展	3
2.1.1 土壤氮素渗漏淋失形态和迁移机理	3
2.1.2 氮素渗漏淋失的研究方法	4
2.1.3 影响土壤氮素渗漏淋失的因素	5
2.1.4 模型研究进展	12
2.2 农田地表径流氮磷流失研究进展	15
2.2.1 氮磷流失与农田耕作状况	15
2.2.2 氮磷流失与降雨	16
2.2.3 氮磷流失与土地利用	17
2.2.4 氮磷流失与肥料施用	18
2.2.5 氮磷流失与下垫面性质	19
2.2.6 氮磷流失与植被覆盖	19
2.2.7 氮磷流失与作物生长季节的关系	21
2.3 氮同位素技术在面源氮污染研究中的应用进展	21
2.3.1 测试技术及方法	21
2.3.2 氮同位素在流域水体硝酸盐污染研究中的应用	23
2.4 氮在溪流中的迁移和转化	25
2.4.1 溪流沿程 $\text{NH}_4\text{-N}$ 和 $\text{NO}_3\text{-N}$ 的变化特征	25
2.4.2 氮在溪流中的硝化作用	25
2.4.3 氮在溪流中的反硝化作用	26
2.4.4 氮在溪流中的生物吸收过程	26
3 研究目标、内容和技术路线	27
3.1 研究目标	27
3.2 研究内容	27

3.3 研究技术路线	27
4 研究区概况与实验方案	29
4.1 研究区概况	29
4.1.1 地理位置与地貌	29
4.1.2 气象条件	30
4.1.3 土壤和植被条件	30
4.1.4 农业种植状况	31
4.1.5 试验小流域概况	32
4.2 实验方案	33
4.2.1 土壤渗漏水监测实验设计	33
4.2.2 浅层地下水监测实验设计	35
4.2.3 地表径流监测实验设计	36
4.2.4 硝酸盐 $\delta^{15}\text{N}$ 同位素的测定	37
4.2.5 微环境实验	39
5 五川流域氮素流失过程与机理研究	41
5.1 地表径流输出	41
5.1.1 降雨径流	41
5.1.2 基流	50
5.1.3 地表径流	60
5.2 地下渗漏淋失	62
5.2.1 土壤剖面氮素分布情况	62
5.2.2 氮在渗漏水与浅层地下水的形态分布	65
5.2.3 浅层地下水氮的浓度与形态特征	67
5.2.4 地下水浓度的季节变化	69
5.2.5 影响氮渗漏淋失的因素	70
5.2.6 氮素渗漏淋失负荷	72
5.3 结论	83
6 五川流域溪流氮的转化过程	85
6.1 溪流沿程 $\text{NO}_3\text{-N}$ 和 $\text{NH}_4\text{-N}$ 的变化特征	86
6.1.1 浓度变化特征	86
6.1.2 原因分析	87
6.2 流域地表径流氮素输入	88

6.3 地下潜流补给	88
6.4 溪流的硝化作用	89
6.4.1 溪流源头和出口 $\text{NO}_3\text{-N}$ 和 $\delta^{15}\text{N}$ 的变化特征	89
6.4.2 溪流沿程 $\text{NO}_3\text{-N}$ 、 $\text{NO}_2\text{-N}$ 、 DO 和 $\delta^{15}\text{N}$ 的变化	90
6.4.3 溪流表层水体 $\text{NO}_3\text{-N}$ 与 $\text{NH}_4\text{-N}$ 动态变化	96
6.5 溪流的反硝化作用	98
6.6 溪流的生物吸收作用	98
6.7 讨论	98
6.8 结论	98
7 总结	101
7.1 主要研究结论	101
7.1.1 五川流域地表径流输出过程及机制	101
7.1.2 五川流域氮的渗漏淋失过程及机制	102
7.1.3 五川流域溪流氮的迁移和转化	103
7.2 主要创新点	103
7.3 不足之处	104
7.4 研究展望	104
参考文献	105
附 录	117
附录 1 参加课题和学术会议	117
附录 2 研究生期间发表论文和所获奖项	118
致 谢	119

厦门大学博硕士论文摘要库

CONTENTS

LIST OF TABLES	IX
LIST OF FIGURES	XI
1 PREFACE.....	1
2 REVIEW OF LITERATURE	3
2.1 Nitrogen leaching in agriculture fields	3
2.1.1 Mechanisms of nitrogen leaching	3
2.1.2 Research method.....	4
2.1.3 Factors of influencing nitrogen leaching	5
2.1.4 Progress of model development.....	12
2.2 Nitrogen and phosphorus discharge of stream runoff.....	15
2.2.1 Impact of tillage	15
2.2.2 Impact of rainfall	16
2.2.3 Impact of landuse.....	17
2.2.4 Impact of fertilizer and manure application.....	18
2.2.5 Impact of underlaying surface	19
2.2.6 Impact of vegetation cover.....	19
2.2.7 Impact of crop growth season.....	21
2.3 Application of nitrogen isotope technique on non-point pollution.....	21
2.3.1 Approach and merhod.....	21
2.3.2 Application of nitrogen isotope tracing method in nitrate pollution-of catchment ...	23
2.4 Nitrogen transformations in stream	25
2.4.1 Variation of $\text{NH}_4\text{-N}$ and $\text{NO}_3\text{-N}$ with downstream.....	25
2.4.2 Nitrification in stream	25
2.4.3 Denitrification in stream	26
2.4.4 Biological assimilation in stream.....	26
3 OBJECTIVES AND APPROACH OF STUDY	27
3.1 Objectives	27
3.2 Contents.....	27

3.3 Approach.....	27
4 STUDY SITE DESCRIPTION AND EXPERIMENT DESIGN....	29
4.1 Description of study site.....	29
4.1.1 Geographical condition.....	29
4.1.2 Meteorological condition.....	30
4.1.3 Soil and vegetation condition.....	30
4.1.4 Agriculture	31
4.1.5 Experimental site	32
4.2 Experiment design and methods.....	33
4.2.1 Experiment design of monitoring nitrogen leaching.....	33
4.2.2 Experiment design of monitoring shallow groundwater.....	35
4.2.3 Experiment design of monitoring streamflow.....	36
4.2.4 Method of testing $\delta^{15}\text{N}$ of nitrate in surface water.....	37
4.2.5 Microcosms experiments	39
5 NITROGEN LOSS OF WUCHUAN CATCHMENT	41
5.1 Stream distrage of nitrogen.....	41
5.1.1 Stormflow	41
5.1.2 Baseflow	50
5.1.3 Nitrogen load in streamflow	60
5.2 Nitrogen leaching	62
5.2.1 Nitrogen distribute of soil section.....	62
5.2.2 Distribute of different nitrogen forms in percolation water and groundwater	65
5.2.3 Nitrogen concentration and forms in shanllow groundwater.....	67
5.2.4 Seasonal variation of nitrogen concentration in groundwater.....	69
5.2.5 Factors of influencing nitrogen leaching	70
5.2.6 Nitrogen leaching load.....	72
5.3 conclusions	83
6 NITROGEN TRANSFORMATION IN WUCHUAN STREAM ..	85
6.1 Variation of $\text{NH}_4\text{-N}$ and $\text{NO}_3\text{-N}$ with downstream	86
6.1.1 Variation of nitrogen concentration.....	86
6.1.2 Casues and consequences	87
6.2 Nitrogen imports of streamflow	88

Degree papers are in the "[Xiamen University Electronic Theses and Dissertations Database](#)". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.

厦门大学博硕士论文摘要库